

# On the interpretation of the latest AMS-02 cosmic ray electron spectrum

In collaboration with M. Di Mauro & S. Manconi  
Based on <https://arxiv.org/abs/2010.13825> (sub. PRD)

Fiorenza Donato  
Torino University

ICRC 2021 - Berlin (online)  
July 15, 2021

# Sources of cosmic $e^-$ and $e^+$ in the Milky Way

Our goal is to reproduce the spectrum of  $e^-$  spectrum measured by

AMS-02 (PRL, 122 (2019) 101101)

Simultaneously, we fit the  $e^+$  AMS-02 data (PRL, 122 (2019) 041102)




Assess the nature of the hardening in the  $e^-$  data around 42 GeV.

1. Secondary  $e^-$ ,  $e^+$
2. Primary  $e^+e^-$  from pulsar
3. Primary  $e^-$  from SNR

- Smooth (Green 2015) distribution of sources
- Diffusion and energy losses (Sync, Inverse Compton scattering (ICS))
- $Q(E)_{\text{SNR}}$ : power law with expo cutoff,  $Q(E)_{\text{PWN}}$ : broken power law

Focus on energy losses due to ICS


# Energy loss rate on the ISRF



Changing the ISRF from full (Vernetto & Lipari 2016) to black body approx is not relevant.

Relevant is changing full ICS computation (numerical) with approximation by (Schlickeiser & Ruppel 2016), applied in (Evoli, Blasi, Amato, Aloisio PRL2020). ICS approximation is wrong for AMS-02 energies (Fang, Bi, Lin arXiv:2007.15601). It induces visible change of slope (starlight - CMB)

# Effect of energy losses on the $e^-$ flux



Different treatments for ICS losses implemented in  $e^-$  flux computation

- ⊕ ICS approximated cross section →  $e^-$  flux following AMS-02 data
- ⊕ Thomson approximation too soft
- ⊕ Full numerical ICS → no evident slope change

Another option to explain AMS-02 data?

# Positrons, and the break in the $e^-$ spectrum

$e^+$  and  $e^-$  AMS-02 spectra FITTED with a multi-component model

$$Y_{1,PWN}=1.88, Y_{1,PWN}=2.31, \eta_{PWN}=0.91\%, Y_{SNR}=2.57, W_{SNR}=1.4 \cdot 10^{49} \text{ erg}, q_{sec}=1.32$$










FIG. 4. Left Panel: Result for the combined fit to  $e^-$  and  $e^+$  AMS-02 data (black and grey data points). We show the secondary production of  $e^+$  (dashed green line) and  $e^-$  (dotted green line),  $e^\pm$  from PWNe (solid red line),  $e^-$  from SNRs (dot-dashed blue line). Right Panel: same as the left panel but zooming in the  $e^-$  sector.

Full (numerically) energy losses kept into account  
The break at 42 GeV is well explained by the interplay  
between SNR and PWN contribution

# Different inputs: impact on e- flux



Changes in the ISRF and/or propagation parameters irrelevant

ICS (wrong) approximation shapes SNR e- flux

Significance for PWN contribution is 4-8%

## Conclusions

- We have demonstrated that approximated ICS cross section gives a bad description in AMS-02 energy range
- Within this approximation, we recover AMS-02 slope change at 42 GeV
- Full numerical ICS does not predict  $e^-$  slope change
- AMS-02  $e^-$  and  $e^+$  data are naturally fitted with dominant SNR  $e^-$  and  $e^\pm$  from PWNe
- The break measured by AMS-02 in the  $e^-$  flux at  $\sim 40$  GeV is very likely due to the interplay between SNR and PWN emission