

Suppression of the TeV pair-beam plasma instability by a weak intergalactic magnetic field

Mahmoud Alawashra

Martin Pohl

July 2021

University of Potsdam & DESY




Table of contents

1. Introduction
2. Weak Intergalactic Magnetic Fields effect on the Linear Growth Rate of Electrostatic Instability
3. Plasma instability limit
4. Summary

Introduction

Introduction


- Blazars are AGN's with a jet oriented along the line of sight.
- Some population of blazars (BL Lacs, in particular) shows an intense emission γ -ray at TeV energies.
- Along with the primary TeV emission we expected to detect an electromagnetic cascade in the GeV energy band due to the attenuation in the IGM:



The electromagnetic cascade is missing in the observations


- Some of the observed blazars arriving energy fluxes in the GeV band are under the predicted flux from the full electromagnetic cascade.

Fig. 1. A comparison of models of cascade emission from TeV blazars (thick solid black curves) with Fermi upper limits (gray curves) and HESS data (gray data points). Thin dashed curves show the primary (unabsorbed) source spectra. Dotted curves show the overall (cascade plus direct) spectra after propagation through the EBL. Vertical lines with arrows show the energies below which the cascade emission should be suppressed. HESS data points (with SEM error bars) are taken from (23, 36, 37).




Two possible explanations

Deflection by the IGM magnetic field




Neronov and Vovk (2010)
Taylor et al. (2011)

Energy loss due to the
Beam-plasma instabilities



Broderick et al. (2012)
Brejzman and Ryutov (1974)

What is the effect of weak intergalactic magnetic fields on the plasma instability?



**Weak Intergalactic Magnetic
Fields effect on the Linear
Growth Rate of Electrostatic
Instability**

Weak Intergalactic Magnetic Fields effect on the Linear Growth Rate of Electrostatic Instability


- The linear growth rate depends significantly on the pair beam distribution.
- The intergalactic magnetic fields cause stochastic deflections of the electrons and positrons increasing the angular distribution function of the pair beam as a Gaussian with the angle spread

$$\Delta\theta = \frac{m_e c}{p} \left[1 + \sqrt{\frac{2}{j} \lambda_{IC} \lambda_B} \frac{e B_{IGM}}{m_e c} \right]$$

$$\approx \approx \lambda_B = \mu c \quad 10^{-6} \left[1 + \frac{B_{IGM}}{10^{-18} [G]} \right]$$

$$w_i \propto \frac{1}{(\Delta\theta)^2}$$

Maximum linear growth rate with IGMF



Plasma instability limit

Energy loss time of the beam-plasma instability

- We consider the energy loss time in Vafin et al. (2018) that is about one order of magnitude less than the IC energy loss time at redshift 0.2:


$$\frac{\tau_{\text{loss}}}{\tau_{\text{IC}}} \approx 0.15,$$

- The energy loss time of the beam-plasma instability is inversely proportional to the maximum linear growth rate (Miniati and Elyiv, 2013; Vafin et al., 2018)

$$\tau_{\text{loss}} \propto \omega_{i,\text{max}}^{-1}.$$

- → The weak intergalactic magnetic field increases the energy loss time of the beam-plasma instability suppressing it after a certain limit.

Plasma instability limit compared to the time delay limit



Summary

Summary

- Weak intergalactic magnetic field slow down the linear electrostatic instability.
- This suppression is effective for fields a factor of a thousand weaker than those needed for magnetic deflection of the cascade emission.
- Non-observation of the cascade emission hence exclude IGM magnetic field of intermediate strength (10^{-15} G on parsec scale).

References

- B.N. Brejzman and D.D. Ryutov. Powerful relativistic electron beams in a plasma and in a vacuum (theory). *Nuclear Fusion*, 14(6):873–907, dec 1974. doi: 10.1088/0029-5515/14/6/012. URL <https://doi.org/10.1088/0029-5515/14/6/012>.
- Avery E. Broderick, Philip Chang, and Christoph Pfrommer. THE COSMOLOGICAL IMPACT OF LUMINOUS TeV BLAZARS. i. IMPLICATIONS OF PLASMA INSTABILITIES FOR THE INTERGALACTIC MAGNETIC FIELD AND EXTRAGALACTIC GAMMA-RAY BACKGROUND. *The Astrophysical Journal*, 752(1):22, may 2012. doi: 10.1088/0004-637x/752/1/22. URL <https://doi.org/10.1088/0004-637x/752/1/22>.
- Francesco Miniati and Andrii Elyiv. RELAXATION OF BLAZAR-INDUCED PAIR BEAMS IN COSMIC VOIDS. *The Astrophysical Journal*, 770(1):54, may 2013. doi: 10.1088/0004-637x/770/1/54. URL <https://doi.org/10.1088/0004-637x/770/1/54>.
- A. Neronov and I. Vovk. Evidence for strong extragalactic magnetic fields from fermi observations of tev blazars. *Science*, 328(5974):73–75, Apr 2010. ISSN 1095-9203. doi: 10.1126/science.1184192. URL <http://dx.doi.org/10.1126/science.1184192>.
- A. M. Taylor, I. Vovk, and A. Neronov. Extragalactic magnetic fields constraints from simultaneous GeV-TeV observations of blazars. , 529:A144, May 2011. doi: 10.1051/0004-6361/201116441.

S. Vafin, I. Rafighi, M. Pohl, and J. Niemiec. The electrostatic instability for realistic pair distributions in blazar/ebl cascades. *The Astrophysical Journal*, 857(1):43, Apr 2018. ISSN 1538-4357. doi: 10.3847/1538-4357/aab552. URL <http://dx.doi.org/10.3847/1538-4357/aab552>.